

Weather Conditions in Southern Africa for Long Distance and High Speed Soaring Flights or The "Magic" of the Trough Line

by **Helmuth Fisher**

Introduction

Extraordinary soaring flights, such as Klaus Holighaus' world record over a 1400 km triangular course and Helmuth Fischer's speed world record over a 1000 km triangular course of 169,7 km/h, have put the spotlight on South Africa as one of the prime "hunting grounds" for world records.

Due to the lack of understanding of Southern Africa's weather mechanism, the potential of this area has not been fully explored. Record attempts were mainly undertaken during relatively short periods and launch points were not systematically chosen. This is not surprising as to date no in-depth analysis exists of the South African weather conditions in respect of soaring flights.

With this paper the author attempts to shed some light on the subject.

The Weather Pattern of the South African Summer

Most of the interior of South Africa is situated at an elevation of between 3.000 and 5.000 ft MSL. Therefore, only atmospheric conditions above 950 - 850 mb are of interest for the soaring areas under discussion. During the summer months, the interior is screened off from the effect of frontal systems by coastal mountain ranges, in particular to the South of the country.

Two oceans surround the "Southern Cape" of Africa. The Atlantic Ocean to the West and the Indian Ocean to the East and South East. Two currents determine the temperature of the two oceans. Originating from the Antarctic region, the cold Benguela current brushes the Western coast of the country. In contrast a warm current emanating from an area South of Madagascar affects the Indian Ocean to the East and the South East of the country.

The major weather engine for the Southern African region is a strong high pressure system over the Atlantic Ocean, which periodically sweeps around the country to link up with a high pressure system over the Indian Ocean. The outstanding and, for soaring, most important feature in this "concert of air masses" is the formation of a heat low over the interior of Southern Africa during the summer months. Due to its elongated shape this is frequently referred to as the "trough line".

[Figures 1](#) and [Figures 2](#) show the typical summer weather pattern of Southern Africa at the 850 mb level and at 13.000 ft respectively. The corresponding weather pattern is as follows:




Fig 1 - 850mb - Typical Summer Weather Pattern of SA




Fig 2 - 13 000ft - Typical Summer Weather Pattern of SA

Warm moist air is moving in from the East. Dewpoints in the area around Johannesburg would be typically in the region of 15 - 17 degrees Celsius. The inversion East of the trough line is predominantly weak or non-existent, which frequently leads to the development of thunderstorms. Therefore, the interior of South Africa is also referred to as the summer rainfall region.

[Figure 3](#) shows the lightning density that supports the abovementioned statement. For example, Johannesburg has one of the highest lightning densities in the world.




Fig 3 - Lightning ground flash density (1975-1986)

Having lost most of its moisture, the subsiding air to the West of the trough line, together with a strong high pressure system in the upper air, produces cumulus clouds with the characteristically high cloud base for which South Africa and Namibia are well known in gliding circles.




Fig 4 - Summer Rainfall Dec\Jan\Feb - 1921-1950

[Figure 4](#) shows the average rainfall in Southern Africa. It can be seen how the trough line separates arid regions to the West from prosperous agricultural land in the East. Maize and cattle farming characterise the area East of the trough line whereas sheep farming dominates the area West of the trough line.

The Significance of the Trough Line for Soaring.

The influx of cool unstable air, which in central Europe during spring can lead to extraordinary soaring conditions, is unknown in South Africa. Only convective processes as a result of high temperatures play a role. Cool Antarctic air pushed into the interior from the South or South-West by the Atlantic High, is an absolute "killer" for soaring conditions.

As a result of regular thunderstorms the high moisture content of the soil East of the trough line has a similar detrimental effect. This, together with the high dewpoints, leads to a low cloud base (4.000 to 5.000 ft AGL) and weak thermal conditions in the East, e.g. Johannesburg area.

This situation is vastly improved closer to the trough line where due to moisture loss of the air the cloud base is lifting. However dewpoints are still high enough and together with a weak inversion, thunderstorm development prevails. Subsiding air just West of the trough line forms an inversion - so vitally important for successful soaring flights. This subdues the thunderstorm activity and together with the now significantly dryer air (dewpoints near De Aar between 3 and 7 degrees Celsius), leads to the formation of flat cumulus clouds with a cloud base of between 14.000 and 18.000 ft MSL as well as strong thermal activity - a glider pilot's dream.

[Figure 5](#) and [Figure 6](#) show typical summer tephigrams of the Johannesburg area and just West of the trough line (De Aar) respectively which illustrate the above effects.




Fig 5 - Tephigram 00.00 GMT for De Aar




Fig 6 - Tephigram 00.00 GMT for Johannesburg

Even further West conditions for soaring deteriorate with the Atlantic High now coming into effect with more stable air and a drop in the inversion level.

Therefore, the position of the Atlantic High, but more importantly, the position and the vertical extension of the heat low (trough line) and their movements during the course of the day, become the critical planning parameters not only for the setting of the task, but also for the selection of the launch point.

The cyclical movements of the Atlantic High around the country linked to the passage of cold fronts South of the Cape Peninsula determine the position of the trough line, which oscillates around a clearly defined mean. This is a line stretching from Grootfontein in Namibia to almost East London in South Africa. It is illustrated in [Figures 7](#), [Figure8](#), [Figure9](#) and [Figure 10](#). Soaring conditions are thus becoming predictable and, depending on the choice of the launch point, a high flexibility of task setting can be obtained.




Fig 7 - Typical Trough Line Movement - During Summer




Fig 8 - Typical Trough Line Movement - During Summer




Fig 9 - Typical Trough Line Movement - During Summer




Fig 10 - Typical Trough Line Movement - During Summer

Given a location, e.g. De Aar, which most of the time is situated just West of the trough line, [Figure 11](#) shows the effect of the approaching trough or the deepening of the heat low. The latter two days of this sequence provided conditions in which two outstanding world records were flown. [Figures 12](#), [Figure 13](#) & [Figure 14](#) show the related tephigrams of De Aar (DY), Upington (UP) and Bloemfontein (BL), on one of the days, namely 5 January 1995.




Fig 11- Rising inversion level as a result of the intensifying of the heat low




Fig 12 - Tephigram 05.01.95 - 00.00 - De Aar






Fig 13 - Tephigram 05.01.95 - 00.00 - Upington



The author used this day to fly a speed world record over a 1.000 km triangular course with an average speed of 169,7 km/h. Judging from the general conditions in the task area, it was estimated that a speed of approximately 155 km/h could have been achieved over a triangular course of 1.250 km. In [Figure 15](#) the aforementioned is placed in relation to the world records existent at that time. The comparison points to the tremendous potential available in South Africa. The high altitude at which flights are performed definitely leads to high speeds.



Timing and Location for Record Flights in South Africa

Being entirely dependent on the weather, there is never an absolute guarantee linked to a certain place. Therefore, the choice is aimed at increasing the probability for success. From the above deliberations it can be gleaned that a location on or just West of the average position of the trough line represents an optimum.

[Figure 16](#) depicts the area in question.




Fig 16 - Recommended area for high speed and long distance soaring flights

In this paper no consideration will be given to other criteria, such as outlanding facilities or luxurious accommodation. These would lead to the choice of sub-optimum locations, such as Kimberley, Bloemfontein or Mafikeng (previously Mmabatho), all of which are located East of the trough line.

Irrespective of the above, the availability of adequate accommodation and a suitable airfield play an important role when selecting a launch point. Considering all criteria, most points are scored by **Gariep Dam (previously Hendrik Verwoerd Dam)**. The most spectacular world records flown in South Africa in recent years confirm this.

[Figure 17](#) shows what a typical triangular task flown from **Gariep Dam** would look like. The launch point would be situated on one leg of the triangle. Therefore the first short leg would lead to the East utilising the prevailing tailwind associated with the high-pressure system in the upper air (see [Figure 2](#)). On the second leg this would turn into a headwind, which at times can be rather strong, but would reduce further to the North-West as one approaches the centre of the upper air High. On the second, and more so on the last leg, one would again benefit from the tailwind. This is particularly welcome towards the evening when returning from a long distance task.




Fig 17 - Typical triangular task flowing from GARIEP DAM 03.00 GMT - 05.01.95

The synoptic situation, which would provide the desired properties, prevails during the months of November up to and including February. In some years even March can provide surprisingly good conditions, albeit at reduced sunshine hours available. The author has experienced a cloud base of 16.000 ft with climb rates of between 5 - 6 m/s during March.

Soaring Conditions in Countries North of South Africa

The summer circulation also benefits a place such as Bitterwasser in Namibia. Many excellent records flown there over the years confirm this.

However, some constraints influence soaring conditions around Bitterwasser. As a result of the summer circulation, moist subtropical air is moving South in the upper air from the Inter-tropical Convergence Zone. This process intensifies as the summer progresses and brings extensive rain to Namibia from approximately the middle of January through to March. This not only limits the number of months during which records can be flown, but at times limits the task area in the North of Bitterwasser due to substantial Altostratus and Altocumulus development.

The other constraint is the result of the sea breeze. In Southern Africa, particularly in the Western parts, the sea breeze can at times move over a distance of up to 300 km inland. Originating from the cold Atlantic Ocean off the West coast of Namibia the effect on soaring conditions can be disastrous. If the stable sea breeze air is not pushed back during the night, thermal development can still be delayed the following morning.

The Northern part of Namibia, Southern Angola, Botswana, Zambia and Zimbabwe are fully affected by the influx of moist sub-tropical air during the summer and, therefore, these areas are undesirable for record attempts during the months November to March. The best months for these countries are September/October and April/May. However, available daylight hours are already reduced during these months. [Figures 18](#) and [Figure 19](#) indicate the approximate suitable areas for soaring flights during the months in question. September and October - being drier - are the better months for record attempts.




Fig 18 - Sept/Oct - Recommended excellent soaring conditions North of South Africa




Fig 19 - April / May - Recommended excellent soaring conditions North of South Africa

The constraints for these areas are the accessibility, the limited number of suitable airfields and the very limited number of places with adequate accommodation and, most importantly, the total lack of outlanding facilities. The latter, as well as logistical difficulties necessitate the exclusive use of motor-gliders for record attempts. One of the hazards which may be encountered during an outlanding is the fact that parts of the flying area is still inhabited by wild animals, such as lions.

The author has flown in some of the areas further North and did not encounter insurmountable administrative difficulties. This is different in Zimbabwe and may be problematic in Southern Angola, which is still in the process of recovering from a civil war.

In the past a major difficulty has been the lack of accurate map material, as well as the lack of suitable turnpoints. GPS assisted navigation, and more recently, documentation using GPS data loggers by which turnpoints can simply be defined by their co-ordinates has basically eliminated those problems.